

Phenolic Compounds from *Scorzonera tomentosa* L.

by Aynur Sanı^{a)}, Christian Zidorn^{b)}, Ernst P. Ellmerer^{c)}, Fevzi Özgökçe^{d)}, Karl-Hans Ongania^{c)}, and Hermann Stuppner^{b)}

^{a)} Istanbul University, Faculty of Pharmacy, Department of Pharmacognosy, 34 116 Beyazıt, Istanbul, Turkey (phone: +90-212-4400000/13581; fax: +90-212-4400252; e-mail: aynur@istanbul.edu.tr)

^{b)} Institut für Pharmazie der Universität Innsbruck, Pharmakognosie, Innrain 52, 6020 Innsbruck, Austria

^{c)} Institut für Organische Chemie der Universität Innsbruck, Innrain 52a, 6020 Innsbruck, Austria

^{d)} Yüzüncü Yıl University, Faculty of Sciences and Letters, Department of Biology, 65180 Van, Turkey


From the subaerial parts of *S. tomentosa* L. (Asteraceae), two new dihydroisocoumarins, compounds **1** and **2**, a new phthalide, **3**, and a new stilbene derivative, **4**, were isolated, together with four known compounds, (±)-hydrangenol (**5**), (–)-hydrangenol 4'-O-β-glucoside (**6**), (±)-hydramacrophylloside A (**7**), and (±)-hydramacrophylloside B (**8**). All secondary metabolites were identified on the basis of physicochemical, spectroscopic, and mass-spectrometric data. The known compounds **5**–**8** were isolated for the first time from this species.

Introduction. – The genus *Scorzonera* is a member of the family Asteraceae, which is a known source of numerous classes of bioactive natural products [1]. For the European flora, some 28 *Scorzonera* species have been reported [2], and the flora of Turkey encompasses 39 species, including 17 endemics [3]. Previous chemical investigations of this genus yielded dihydroisocoumarins, flavonoids, lignans, phenolic acids, a sesquiterpene, sesquiterpene lactones, triterpenes, and a new class of bibenzyl derivatives [4–17].

Scorzonera tomentosa L. is a perennial herb endemic to Turkey [3]. Its subaerial parts are used in traditional medicine as analgesic, antirheumatic (in the form of plaster), and anthelmintic, as well as for the treatment of infertility [18] [19]. In a previous paper [17], the isolation of stigmasteryl 3β-glucoside, β-sitosterol, lupeol, lupeol acetate, and α-amyrin from the aerial parts of *S. tomentosa* was reported. In the current study, we report four new constituents, compounds **1**–**4**, from the subaerial parts of this plant, together with four known compounds: (±)-hydrangenol (**5**), (–)-hydrangenol 4'-O-glucoside (**6**), and (±)-hydramacrophylloside A (**7**) and B (**8**).

Results and Discussion. – The AcOEt-soluble fraction of the MeOH extract of the subaerial parts of *S. tomentosa* L. was investigated to elucidate its secondary-metabolite profile.

Compound **1** was isolated as a colorless, optically inactive powder ($[\alpha]_D^{25}=0$). The UV spectrum of **1** showed absorption maxima at 246 (log $\epsilon=4.1$) and 306 (3.9) nm. The IR spectrum showed absorption bands due to phenolic OH (3450), C=O (1736), and aliphatic groups (3063 cm⁻¹), as well as due to an aromatic ring (1600, 1584, 1475 cm⁻¹). In the ESI mass spectrum, the $[M-H]^-$ signal was observed at m/z 269, consis-



tent with the molecular formula C₁₆H₁₄O₄. The ¹H-NMR spectrum of **1** (Table 1) showed the signals ascribable to a 1,4-disubstituted benzene ring [δ (H) 7.32 (*d*, *J* = 8.5 Hz, 2 H); 6.81 (*d*, *J* = 8.5 Hz, 2 H)], a trisubstituted benzene ring [δ (H) 6.94 (*d*, *J* = 7.5 Hz, 1 H); 7.56 (*t*, *J* = 7.5 Hz, 1 H); 7.08 (*d*, *J* = 8.5 Hz, 1 H)], a δ -lactone [δ (H) 5.40 (*dd*, *J* = 3.0, 12.0 Hz, 1 H); 3.03 (*dd*, *J* = 3.0, 16.0 Hz, 1 H); 3.26–3.36 (*m*, 1 H)], and a MeO group [δ (H) 3.91 (*s*, 3 H)].

The ¹³C-NMR spectrum of **1** (Table 2) revealed the presence of twelve aromatic C-atoms (three quaternary, two oxygenated quaternary, and seven CH), as well as one CH₂, one oxygenated CH, one MeO, and one C=O group. The position of the MeO group was established at C(8) from the ¹H-NMR coupling pattern, and corroborated by HMBC experiments (Table 3). In the HMBC spectrum, H–C(5) showed a strong three-bond correlation with C(4) at δ (C) 37.05, and the MeO H-atoms at δ (H) 3.91 were correlated with C(8) at δ (C) 162.6. From these data, the structure of racemic **1** was established as (3*RS*)-3,4-dihydro-3-(4-hydroxyphenyl)-8-methoxy-1*H*-2-benzopyran-1-one, and named (±)-*scorzotomentosin*.

Compound **2** was isolated as a colorless powder. HR-FAB-MS displayed signals at *m/z* 455.1325 ([*M* + Na]⁺; calc. 455.1318) and 433.1507 ([*M* + H]⁺; calc. 433.1499), consistent with the molecular formula C₂₂H₂₄O₉. The data observed in the ¹H- and ¹³C-NMR spectra (Tables 1 and 2, resp.) were very similar to those of **1**, but further indicated the presence of a sugar moiety. The anomeric H-atom at δ (H) 4.94 (*d*, *J* = 7.0 Hz) and the coupling pattern of the other sugar resonances indicated a β -glucose (Glc) unit. The position of the glucoside linkage was established by a HMBC experiment (Table 3), which revealed correlations between the anomeric H-atom, H–

Table 1. ^1H -NMR Spectroscopic Data of **1–4**. At 500 MHz in CD_3OD (**1–3**) or $(\text{D}_6)\text{DMSO}$ (**4**); δ in ppm, J in Hz. Arbitrary atom numbering.

Position	1	2 ^{a)}	3	4
3	5.40 (<i>dd</i> , $J=3.0$, 12.0)	5.46 (<i>dd</i> , $J=3.0$, 12.0)	5.63 (<i>d</i> , $J=4.0$)	–
4	3.03 (<i>dd</i> , $J=3.0$, 16.0)	3.17 (<i>dd</i> , $J=3.0$, 16.0)	6.73 (<i>d</i> , $J=8.0$)	6.77 (<i>d</i> , $J=8.0$)
	3.26–3.36 (<i>m</i>)	3.26–3.36 (<i>m</i>)		
5	6.94 (<i>d</i> , $J=7.5$)	6.94 (<i>d</i> , $J=7.5$)	7.53 (<i>dd</i> , $J=7.5$)	7.10 (<i>dd</i> , $J=7.5$, 8.0)
6	7.56 (<i>t</i> , $J=7.5$)	7.56 (<i>t</i> , $J=8.5$)	7.00 (<i>d</i> , $J=8.5$)	7.26 (<i>d</i> , $J=8.0$)
7	7.08 (<i>d</i> , $J=8.5$)	7.08 (<i>d</i> , $J=8.5$)	–	–
8	–	–	5.03 (<i>d</i> , $J=4.0$)	–
2',6'	7.32 (<i>d</i> , $J=8.5$)	7.44 (<i>d</i> , $J=8.5$)	7.12 (<i>d</i> , $J=8.5$)	7.34 (<i>d</i> , $J=8.5$)
3',5'	6.81 (<i>d</i> , $J=8.5$)	7.14 (<i>d</i> , $J=8.5$)	6.68 (<i>d</i> , $J=8.5$)	6.82 (<i>d</i> , $J=8.5$)
α	–	–	–	7.13 (<i>d</i> , $J=16.0$)
β	–	–	–	7.03 (<i>d</i> , $J=16.0$)
MeO	3.91 (<i>s</i>)	3.91 (<i>s</i>)	3.91 (<i>s</i>)	3.71 (<i>s</i>)

^{a)} Sugar resonances: $\delta(\text{H})$ 4.94 (*d*, $J=7.0$, $\text{H}-\text{C}(1'')$); 3.48 (*dd*, $J=9.0$, 7.0, $\text{H}-\text{C}(2'')$); 3.45 (*t*, $J=9.0$, $\text{H}-\text{C}(3'')$); 3.40 (*t*, $J=9.0$, $\text{H}-\text{C}(4'')$); 3.42–3.50 (*m*, $\text{H}-\text{C}(5'')$); 3.70 (*dd*, $J=12.0$, 5.0, 1 H of $\text{CH}_2(6'')$); 3.90 (*dd*, $J=12.0$, 2.0, 1 H of $\text{CH}_2(6'')$).

Table 2. ^{13}C -NMR Spectroscopic Data of **1–4**. At 125 MHz in CD_3OD (**1–3**) or $(\text{D}_6)\text{DMSO}$ (**4**); δ in ppm; signal assignment by HSQC and HMBC experiments. Arbitrary atom numbering.

Position	1	2	3	4	Position	1	2	3	4
1	165.5	164.0	170.0	133.1	3'	116.3	117.5	115.0	115.4
2	–	–	–	115.7	4'	159.0	158.0	158.5	158.2
3	81.0	80.5	85.2	154.6	5'	116.3	117.5	115.0	115.4
3a	–	–	150.5	–	6'	128.9	128.4	128.5	127.5
4	37.05	36.0	116.5	109.1	α	–	–	–	123.6
4a	143.9	142.5	–	–	β	–	–	–	128.0
5	120.6	120.5	136.5	126.0	1''	–	102.0	–	–
6	136.4	136.5	112.0	115.7	2''	–	74.6	–	–
7	112.3	112.0	159.0	–	3''	–	77.9	–	–
7a	–	–	115.0	–	4''	–	71.1	–	–
8	162.6	162.0	76.0	–	5''	–	77.7	–	–
8a	114.3	113.5	–	–	6''	–	61.2	–	–
1'	130.8	133.0	129.0	128.0	MeO	56.4	56.2	55.2	54.5
2'	128.9	128.4	128.5	127.5	COOH	–	–	–	172.1

$\text{C}(1'')$, and the quaternary C-atom at $\delta(\text{C})$ 158.0 ($\text{C}(4'')$)¹⁾. Consequently, **2** was identified as (–)-scorzotomentosin 4'-O- β -glucoside.

The absolute configuration at C(3) was determined by comparison of the CD spectrum of **2** with that of scorzocreticin [9]. Compound **2** gave rise to a positive Cotton effect at 241 nm, and a negative one at 263 nm, as in the case of scorzocreticin. This suggested that both compounds have the absolute (*S*)-configuration at C(3).

¹⁾ Arbitrary atom numbering (see chemical formulae).

Table 3. Key HMBC Cross-Peaks for **1** and **2**

H-Atom	C-Atom	
	1	2
H–C(3)	C(1), C(4a), C(2',6')	C(1), C(4a), C(2',6')
H–C(4)	C(4a), C(5), C(8a), C(1')	C(4a), C(5), C(8a), C(1')
H–C(5)	C(4), C(7), C(8a)	C(4), C(7), C(8a)
H–C(6)	C(4a), C(8)	C(4a), C(8)
H–C(7)	C(5), C(8a)	C(5), C(8a)
H–C(2',6')	C(3), C(2',6'), C(4')	C(3), C(2',6'), C(4')
H–C(3',5')	C(1'), C(3',5')	C(1'), C(3'/5')
MeO	C(8)	C(8)
H–C(1'')	–	C(4')

Compound **3**, a colorless, optically inactive powder, showed UV absorbances at 228 (log ϵ = 4.2), 285 (3.6), and 301 (3.7) nm. Its IR spectrum indicated phenolic OH (3440), C=O (1740), and aryl (1620, 1607, 1470 cm^{-1}) groups. The molecular formula $\text{C}_{16}\text{H}_{14}\text{O}_5$ was determined by HR-FAB-MS (m/z 287.0912 ($[M+H]^+$; calc. 287.0919)). The ^1H -NMR spectrum of **3** (Table 1) was similar to that of hydramacrophyllol B (**8**) [20], with a 1,2,3-trisubstituted benzene ring ($\delta(\text{H})$ 6.73 (d , J = 8.0 Hz, 1 H); 7.53 (dd , J = 7.5 Hz, 1 H); 7.00 (d , J = 8.5 Hz, 1 H)), a 1,4-disubstituted aromatic ring [$\delta(\text{H})$ 7.12, 6.68 ($2d$, J = 8.5 Hz each, 2×2 H)]. Furthermore, the signals of a CH adjacent to an OH group, and those of a CH adjacent to a lactone O-atom were observed ($\delta(\text{H})$ 5.03 (d , J = 4.0 Hz, 1 H); 5.63 (d , J = 4.0 Hz, 1 H)). Finally, a MeO group resonated at $\delta(\text{H})$ 3.91 (s).

In the ^{13}C -NMR spectrum of **3** (Table 2), twelve aromatic C-atoms (three quaternary, two oxygenated quaternary, and seven CH), two oxygenated CH groups, one MeO, and one C=O group were observed. The MeO group was placed at C(7), as deduced from the ^1H -NMR coupling pattern and interpretation of the HMBC spectrum (Table 4). In the latter, H–C(4) at $\delta(\text{H})$ 6.73 showed a strong three-bond correlation with C(3) at $\delta(\text{C})$ 85.2, and the MeO H-atoms at $\delta(\text{H})$ 3.91 were correlated with C(7) at $\delta(\text{C})$ 159.0. The configurations at C(3) and C(8) were deduced from the NMR data in comparison to those reported for **8** [20]. Thus, from the above data, the structure of racemic **3** was identified as (3*RS*)-3-[(*SR*)-hydroxy(4-hydroxyphenyl)-methyl]-7-methoxy-2-benzofuran-1(3*H*)-one, and named (\pm)-*scorzophthalide*.

Compound **4** was isolated as a yellowish powder. Its IR spectrum showed absorptions typical of OH (3440), C=O (1660), and benzene-ring moieties (1590, 1515, 1460, 1400 cm^{-1}). The UV spectrum was characteristic for a stilbene derivative [21], with absorption maxima at 217 (log ϵ = 4.2), 306 (4.31), and 320 (4.32) nm. The ESI mass spectrum of **4** displayed the $[M-H]^-$ ion at m/z 269, consistent with the molecular formula $\text{C}_{16}\text{H}_{14}\text{O}_4$. The ^1H -NMR spectrum (Table 1) indicated a 1,4-disubstituted benzene ring, a trisubstituted benzene ring, a MeO group, and two vinylic H-atoms [$\delta(\text{H})$ 7.03, 7.13 ($2d$, J = 16.0 Hz, 1 H each)], which were assigned to an (*E*)-stilbene. The ^{13}C -NMR spectrum (Table 2) indicated 16 C-atoms, including one MeO, nine CH, three quaternary C-atoms, two oxygenated aryl C-atoms, and one COOH group. HMBC experiments (Table 4) showed long-range correlations between the MeO H-atoms at

Table 4. Key HMBC Cross-Peaks for **3** and **4**

H-Atom	C-Atom	
	3	4
H–C(3)	C(1), C(4), C(7a), C(1')	–
H–C(4)	C(3), C(6), C(7a)	C(2), C(6)
H–C(5)	C(3a), C(7)	C(1), C(3), C(4)
H–C(6)	C(4), C(7a)	C(2), C(4), C(α)
H–C(8)	C(3a), C(2',6')	–
H–C(α)	–	C(1), C(2), C(6), C(1')
H–C(β)	–	C(1), C(2',6'), C(α)
H–C(2',6')	C(4'), C(2',6'), C(8)	C(4'), C(2',6'), C(β)
H–C(3',5')	C(1'), C(3',5')	C(1'), C(3',5')
MeO	C(7)	C(3)

δ (H) 3.71 and the quaternary C-atom at δ (C) 154.6 (C(3)), and also between the aromatic resonance at δ (H) 7.10 (H–C(5)) and the two quaternary C-atoms at δ (C) 133.1 (C(1)) and 154.6 (C(3)). Hence, the MeO group was placed at C(3), and the COOH group was attached at C(2). From these data, the structure of **4** was determined as 2-[(*E*)-2-(4-hydroxyphenyl)ethenyl]-6-methoxybenzoic acid, and named *scorzoerzinanin*.

The four known compounds, **5**–**8**, had been isolated before from the genus *Hydrangea*, and were isolated from *S. tomentosa* for the first time. They were identified on the basis of their physicochemical and spectroscopic data [20–24].

A. S. thanks TUBİTAK for a fellowship and the Research Fund of Istanbul University for financial support (fund UDP-694/14032006). The authors wish to thank Dr. Sonja Sturm for recording ESI mass spectra, and Birthe Schubert for CD and ORD measurements.

Experimental Part

General. Column chromatography (CC): silica gel 60 (40–63 μ m; Merck) or Sephadex LH-20 (Sigma-Aldrich). Prep. TLC: precoated silica gel 60 F_{254} plates (0.25 mm; Merck). Semi-prep. HPLC: injection of 40- μ l aliquots of sample soln. (25 mg/ml) on a Merck RP-18 column (LiChrosphere, 250 \times 10 mm, 10 μ m) at a column temp. of 55°; isocratic MeCN/H₂O flow at 2 ml/min; stop time: 45 min, post time: 15 min; detection at 205 nm. UV Spectra: Jasco V-530 spectrophotometer, in MeOH; λ_{\max} (log ϵ) in nm. IR Spectra: Perkin-Elmer 552 spectrophotometer, KBr cells; in cm^{–1}. ¹H- and ¹³C-NMR Spectra: Varian UnityPlus-500 spectrometer, at 500/125 MHz, resp., in CD₃OD or (CD₃)₂SO; δ in ppm rel. to Me₄Si, *J* in Hz. ESI-MS: Finnigan MAT SSQ-7000 mass spectrometer. HR-FAB-MS: Finnigan MAT-95 mass spectrometer, glycerol as matrix; in *m/z*.

Plant Material. *Scorzonera tomentosa* L. was collected from Erzincan, East Anatolia, at an altitude of 1560 m, in July 2004. A voucher specimen (F 12 250) was deposited at the Herbarium of the Faculty of Sciences and Letters, Yüzüncü Yıl University, Turkey.

Extraction and Isolation. The air-dried, ground, subaerial parts of *S. tomentosa* (398 g) were repeatedly extracted with MeOH (7 \times 1 l) at r.t., 4 d each. After solvent evaporation, 60 g of residue was obtained, which was dissolved in MeOH/H₂O 1:2 (300 ml), and then successively extracted with petroleum ether (PE), AcOEt, and BuOH. The AcOEt-soluble part (12 g) was purified by CC (SiO₂; CH₂Cl₂/MeOH 99:1, 98:2, 95:5, 90:10, 80:20, 70:30, 50:50, 0:100; total volume 6 l): 24 fractions (*Fr. 1*–*24*). *Fr. 5* (413 mg) was further separated by CC (Sephadex LH-20; MeOH): seven fractions (*Fr. 5.1*–*5.7*). *Fr.*

5.3 (223 mg) was further separated by prep. TLC (SiO₂; CH₂Cl₂/MeOH 9:1) to afford **1** (86 mg) and **5** (33 mg). *Fr. 11* (495 mg) was purified by CC (*Sephadex LH-20*; acetone): 10 fractions (*Fr. 11.1–11.10*). *Fr. 11.6* (62 mg) was subjected to semi-prep. HPLC (MeCN/H₂O 25:75; isocratic) to afford **3** (3 mg; *t_R* 26.8–30.0). *Fr. 12* (631 mg) was purified by CC (*Sephadex LH-20*; MeOH): eleven fractions (*Fr. 12.1–12.11*). *Fr. 12.7* (241 mg) was further separated by semi-prep. HPLC (MeCN/H₂O 22:78; isocratic) to afford **7** (12.7 mg; *t_R* 21.8–24.8), **8** (17.5 mg; *t_R* 25.9–29.0), and **3** (1 mg; *t_R* 37.5–38.0). *Fr. 17* (1.916 g) was subjected to CC (*Sephadex LH-20*; MeOH) to provide 18 subfractions: *Fr. 17.1–17.18*. *Fr. 17.4*, *Fr. 17.8*, and *Fr. 17.14* contained pure **2** (63 mg), **4** (42 mg), and **6** (8 mg), resp.

(±)-*Scorzotomentosin* (= (3*RS*)-3,4-Dihydro-3-(4-hydroxyphenyl)-8-methoxy-1*H*-2-benzopyran-1-one; **1**). Colorless powder. $[\alpha]_D^{25} = 0$ (*c* = 2.34, MeOH). UV (MeOH): 246 (4.1), 306 (3.9). IR (KBr): 3450, 3063, 1736, 1600, 1584, 1475. ¹H-, ¹³C-, and 2D-NMR: see *Tables 1–3*. ESI-MS: 269 ($[M - H]^-$, C₁₆H₁₃O₄⁻).

(-)-*Scorzotomentosin* 4'-*O*-β-Glucoside (= 4'-[(3*S*)-3,4-Dihydro-8-methoxy-1-oxo-1*H*-2-benzopyran-3-yl]phenyl β-Glucopyranoside; **2**). Colorless powder. $[\alpha]_D^{25} = -142$ (*c* = 2.52, MeOH). CD (MeOH): 241 (pos.), 263 (neg.). ¹H-, ¹³C-, and 2D-NMR: see *Tables 1–3*. HR-FAB-MS: 455.1325 ($[M + Na]^+$, C₂₂H₂₄NaO₈⁺; calc. 455.1318), 433.1507 ($[M + H]^+$, C₂₂H₂₅O₈⁺; calc. 433.1499).

(±)-*Scorzophthalide* (= (3*RS*)-3-[(*SR*)-Hydroxy(4-hydroxyphenyl)methyl]-7-methoxy-2-benzofuran-1(3*H*)-one; **3**). Colorless powder. $[\alpha]_D^{25} = 0$ (*c* = 2.26, MeOH). UV (MeOH): 228 (4.2), 285 (3.6), 301 (3.7). IR (KBr): 3440, 1740, 1620, 1607, 1470. ¹H-, ¹³C-, and 2D-NMR: see *Tables 1, 2*, and *4*. HR-FAB-MS: 287.0912 ($[M + H]^+$, C₁₆H₁₅O₅⁺; calc. 287.0919).

Scorzoerzincanin (= 2-[(*E*)-2-(4-Hydroxyphenyl)ethenyl]-6-methoxybenzoic Acid; **4**). Yellowish powder. UV (MeOH): 217 (4.2), 306 (4.31), 320 (4.32). IR (KBr): 3440, 1660, 1590, 1515, 1460, 1400. ¹H-, ¹³C-, and 2D-NMR: see *Tables 1, 2*, and *4*. ESI-MS: 269 ($[M - H]^-$, C₁₆H₁₃O₄⁻).

REFERENCES

- [1] W. C. Evans, 'Trease and Evans' Pharmacognosy', 14th edn., W. B. Saunders Company, Ltd., London, 1996, p. 50.
- [2] A. O. Chater, in 'Flora Europaea', Eds. T. G. Tutin, V. H. Heywood, N. A. Burges, D. M. Moore, D. H. Valentine, S. M. Walters, D. A. Webb, Cambridge University Press, London, 1976, Vol. 4, pp. 317–322.
- [3] D. F. Chamberlain, in 'Flora of Turkey and the East Aegean Islands', Ed. P. H. Davis, Edinburgh University Press, Edinburgh, 1975, Vol. 5, p. 632.
- [4] C. Zidorn, E. P. Ellmerer-Müller, H. Stuppner, *Helv. Chim. Acta* **2000**, *83*, 2920.
- [5] C. Zidorn, E. P. Ellmerer-Müller, H. Stuppner, *Pharmazie* **2000**, *55*, 550.
- [6] C. Zidorn, R. Spitaler, E. P. Ellmerer-Müller, N. B. Perry, C. Gerhauser, H. Stuppner, *Z. Naturforsch., C* **2002**, *57*, 614.
- [7] C. Zidorn, E. P. Ellmerer-Müller, S. Sturm, H. Stuppner, *Phytochemistry* **2003**, *63*, 61.
- [8] C. Zidorn, B. O. Petersen, V. Udovicic, T. O. Larsen, J. Q. Duus, J. M. Rollinger, K. H. Ongania, E. P. Ellmerer, H. Stuppner, *Tetrahedron Lett.* **2005**, *46*, 1291.
- [9] S. Paraschos, P. Magiatis, E. Kalpoutzakis, C. Harvala, A. L. Skaltsounis, *J. Nat. Prod.* **2001**, *64*, 1585.
- [10] F. Menichini, G. Statti, F. Delle Monache, *Fitoterapia* **1994**, *65*, 555.
- [11] O. V. Bryanskii, V. V. Tolstikhina, A. A. Semenov, *Khim. Prir. Soedin.* **1992**, *28*, 591.
- [12] O. V. Bryanskii, V. V. Tolstikhina, S. V. Zinchenko, A. A. Semenov, *Khim. Prir. Soedin.* **1992**, *28*, 640.
- [13] V. V. Tolstikhina, O. V. Bryanski, A. I. Syrincha, A. A. Semenov, *Khim. Prir. Soedin.* **1988**, *24*, 763.
- [14] V. V. Tolstikhina, A. A. Semenov, *Rastitel' nye Resursy* **1998**, *34*, 77.
- [15] V. V. Tolstikhina, A. A. Semenov, I. A. Ushakov, *Rastitel' nye Resursy* **1999**, *35*, 87.
- [16] J. Li, Q. X. Wu, Y. P. Shi, Y. Zhu, *Chin. Chem. Lett.* **2004**, *15*, 1309.
- [17] S. Öksüz, N. Gören, A. Ulubelen, *Fitoterapia* **1990**, *61*, 92.
- [18] T. Baytop, 'A Dictionary of Vernacular Names of Wild Plants of Turkey', Publication No. 578 of the Turkish Language Society, Turkish History Society Press, Ankara, 1997, p. 52.

- [19] T. Baytop, 'Therapy with Medicinal Plants in Turkey', Publication No. 3255 of the Istanbul University, Sanal Press, Istanbul, 1984, p. 263.
- [20] M. Yoshikawa, H. Shimada, N. Yagi, N. Murakami, H. Shimoda, J. Yamahara, H. Matsuda, *Chem. Pharm. Bull.* **1996**, *44*, 1890.
- [21] T. Hashimoto, M. Tori, Y. Asakawa, *Phytochemistry* **1988**, *27*, 109.
- [22] M. Yoshikawa, E. Uchida, N. Chatani, N. Murakami, J. Yamahara, *Chem. Pharm. Bull.* **1992**, *40*, 3121.
- [23] M. Yoshikawa, E. Uchida, N. Chatani, H. Kobayashi, Y. Naitoh, Y. Okuno, H. Matsuda, J. Yamahara, N. Murakami, *Chem. Pharm. Bull.* **1992**, *40*, 3352.
- [24] M. Yoshikawa, H. Matsuda, H. Shimoda, H. Shimada, E. Harada, Y. Naitoh, A. Miki, J. Yamahara, N. Murakami, *Chem. Pharm. Bull.* **1996**, *44*, 1440.

Received October 6, 2006